



## Estimation of Effect Factors for application to marine eutrophication in LCIA

**Cosme, Nuno Miguel Dias; Hauschild, Michael Zwicky**

*Published in:*  
Abstract book - SETAC Europe 24th Annual Meeting

*Publication date:*  
2014

[Link back to DTU Orbit](#)

*Citation (APA):*  
Cosme, N. M. D., & Hauschild, M. Z. (2014). Estimation of Effect Factors for application to marine eutrophication in LCIA. In *Abstract book - SETAC Europe 24th Annual Meeting* SETAC Europe.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Estimation of Effect Factors for application to marine eutrophication in LCIA

Nuno Cosme and Michael Z. Hauschild

Quantitative Sustainability Assessment (QSA), Department of Management Engineering, Technical University of Denmark (DTU), Kongens Lyngby - Denmark

E-mail contact: [nmdc@dtu.dk](mailto:nmdc@dtu.dk)

---

## 1. Introduction

Marine eutrophication is defined as the set of ecosystem responses to nutrient loadings in the photic zone of marine waters. Ecosystems use the increased availability of nutrients to boost primary production [1]. The eventual decay and degradation of this organic matter by heterotrophic bacteria in bottom waters result in the consumption of dissolved oxygen (DO). These responses turn more significant as they result in impacts to ecosystem quality and local economy, by (i) changing communities' composition and species interaction, enhancing the growth of toxic and harmful algal species, (ii) decreasing water quality, by high turbidity, colour, and smell, hindering water uses, fish production, and aesthetic value, and (iii) depleting DO in bottom waters down to hypoxic or anoxic levels with resulting impacts on the survival of benthic species [2].

In oligotrophic waters, the increase in phytoplankton biomass is mainly consumed by the secondary production rendering little negative impacts from the nutrients enrichment. However, in eutrophic waters, impacts are more frequent and the degradation of the ecosystem quality by excessive depletion of DO causes a significant stress on biota. The impacts of hypoxia on biota depend much on the severity, frequency and duration of the exposure to low DO [3], and both acute and chronic effects can be expected. Exposure to extreme or prolonged hypoxia leads to mass mortalities, but hypoxia also induces many different sub-lethal responses in organisms at the behavioural level (e.g. avoidance strategies), at the physiological level (e.g. limiting growth, activity and respiration rates) and at the ecological level (e.g. altering structure, function, and services of benthic communities) [4]. In general, crustaceans show the highest sensitivity to lower oxygen saturation rates followed by fish, whereas molluscs, cnidarians and priapulids show more tolerance to oxygen depletion [5].

## 2. Materials and methods

A dataset of sensitivities of individual species to hypoxia [5] was used for this study. Relevant species were identified (benthic, demersal, or benthopelagic) and their geographical distribution determined. The selected species were grouped into 5 climate zones - polar, subpolar, temperate, subtropical, and tropical (the sample size and lack of representativeness prevented classification at finer spatial resolutions).

Species Sensitivity Distribution (SSD) curves were produced to estimate the Potentially Affected Fraction of species (PAF) at different levels of DO. The SSD methodology uses a probabilistic model to represent the sensitivity of the community to an environmental stressor through a statistical or empirical distribution function of the sampled sensitivities of the individual biological species [6]. For the application in Life Cycle Impact Assessment (LCIA), the distribution of the sensitivities of individual species to hypoxia is used to estimate the sensitivity to low DO levels of the communities found in each climate zone. The distribution is used to estimate the  $HC50_{EC50}$ , i.e. the concentration of DO (intensity of the stressor) affecting 50% of the species above their  $EC_{50}$  level.

Characterisation Factors (CF) are used in LCIA to convert emissions and resources consumed into impact potentials for specific impact categories. The CF integrates the N-fate term (Fate Factor, FF), the habitat exposure term (Exposure Factor, XF), and the term quantifying the effect (Effect Factor, EF) as shown next:

$$CF[PAF \cdot m^3 \cdot yr / KgN] = FF[yr] \times XF[KgO_2 / KgN] \times EF[PAF \cdot m^3 / KgO_2]$$


The EF expresses the change of effect ( $\Delta PAF$ ) due to a variation of the stressor intensity (depletion of DO, i.e.  $\Delta[O_2]$ ) and it is calculated by:

$$EF = \frac{\Delta PAF}{\Delta[O_2]} = \frac{0.5}{HC50_{EC50}}$$

Where  $HC50_{EC50}$  is either obtained from the SSD method, by calculating  $HC50_{EC50} = 10^{\text{avg}(\log EC_{50})}$  or by calculating the geometric mean of the  $EC_{50}$  data, in accordance to the average gradient method [7].

### 3. Results and discussion

Preliminary results of  $HC50_{EC50}$  were produced for 5 climate zones together with a global default. As an example of the obtained SSD curves and the  $HC50_{EC50}$ , the global case is shown in Figure 1.



**Figure 1. Species Sensitivity Distribution (SSD) curve based on data for sensitivity to hypoxia ( $EC_{50}$ ). Example of the results for the estimation of the global default  $HC50_{EC50}=1.453 \text{ mgO}_2/\text{L}$  ( $n=58$ ). PNAF=Potentially Not Affected Fraction of species.**

The preliminary results for the EF estimation for the 5 climate zones and global default are shown in Table 1.

Climate zone	<i>n</i> taxa	SSD curve fit parameters					$HC50_{EC50}$		EF
		$\alpha$	$\beta$	Slope	Interc.	$R^2$	$\text{mgO}_2/\text{L}$	$\text{kgO}_2/\text{m}^3$	$\text{PAF} \cdot \text{m}^3/\text{kgO}_2$
Polar	13	0.195	0.173	3.188	4.290	0.943	<b>1.568</b>	1.57E-03	<b>318.90</b>
Subpolar	38	0.181	0.555	2.429	4.533	0.967	<b>1.516</b>	1.52E-03	<b>329.88</b>
Temperate	47	0.158	0.593	2.498	4.569	0.966	<b>1.440</b>	1.44E-03	<b>347.30</b>
Subtropical	41	0.237	0.481	2.653	4.340	0.935	<b>1.726</b>	1.73E-03	<b>289.70</b>
Tropical	19	0.242	0.197	3.403	4.154	0.972	<b>1.745</b>	1.75E-03	<b>286.46</b>
Global default	58	0.162	0.630	2.558	4.558	0.978	<b>1.453</b>	1.45E-03	<b>344.10</b>

**Table 1: Results of the calculations delivering the terms of the SSD fit equations and the  $HC50_{EC50}$  used to estimate the Effect Factors (EF) per climate zone, plus a global default EF that includes all 58 species.**

### 4. Conclusions

The spatial differentiation obtained for the EF results was found essential to increase the discriminatory power of the model and to assess the results. This approach to calculate EF will be combined with a suitable methodology for the other factors in the CF expression, i.e. fate of nutrients (FF) and habitat exposure (XF) (in prep.).

### 5. References

- [1] Nixon SW. 1995. Coastal marine eutrophication: A definition, social causes, and future concerns. *Ophelia* 41: 199-219.
- [2] Kelly JR. 2008. Nitrogen in the Environment: Chapter 10. Nitrogen Effects on Coastal Marine Ecosystems. USEPA Paper 50, <http://digitalcommons.unl.edu/usepapapers/50>, accessed 10 May 2012.
- [3] Powers SP, Peterson CH, Christian RR, Sullivan E, Powers MJ, Bishop MJ, Buzzelli CP. 2005. Effects of eutrophication on bottom habitat and prey resources of demersal fishes. *Marine Ecology Progress Series* 302: 233-243.
- [4] Wu RSS. 2002. Hypoxia: from molecular responses to ecosystem responses. *Marine Pollution Bulletin* 45: 35-45.
- [5] Vaquer-Sunyer R, Duarte CM. 2008. Thresholds of hypoxia for marine biodiversity. *Proceedings of the National Academy of Sciences USA* 105: 15452–15457.
- [6] Posthuma L, Suter II GW, Traas TP. 2002. *Species Sensitivity Distributions in Ecotoxicology*. Boca Raton FL: Lewis Publishers, 583 pp.
- [7] Pennington DW, Payet J, Hauschild M. 1994. Aquatic ecotoxicological indicators in life-cycle assessment. *Environmental Toxicology and Chemistry* 23(7): 1796-1807.